

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11) Publication number:

0 455 000 A1

(12)

EUROPEAN PATENT APPLICATION(21) Application number: **91105171.2**(22) Date of filing: **02.04.91**

(51) Int. Cl.⁵: **C07D 401/06, A01N 43/50,
A01N 43/54, A01N 43/78,
A01N 43/40, C07D 417/06,
C07D 405/14, C07D 417/14**

(30) Priority: **03.04.90 JP 87368/90
26.02.91 JP 30815/91**

(43) Date of publication of application:
06.11.91 Bulletin 91/45

(84) Designated Contracting States:
CH DE ES FR GB IT LI NL

(71) Applicant: **mitsui toatsu chemicals, Inc.
2-5 Kasumigaseki 3-chome
Chiyoda-Ku Tokyo 100(JP)**


(72) Inventor: **Kodaka, Kenji
Mitsui Toatsu Miyonodai Daini Aparto 41
2142, Tougou, Mobara-shi, Chiba-ken(JP)
Inventor: Kinoshita, Katsutoshi
Miyonodai Ryo 94, 2142, Tougou
Mobara-shi, Chiba-ken(JP)
Inventor: Nakaya, Michihiko**

**Miyonodai Ryo 26, 2142, Tougou
Mobara-shi, Chiba-ken(JP)
Inventor: Ebihara, Koichi
Miyonodai Ryo 19, 2142, Tougou
Mobara-shi, Chiba-ken(JP)
Inventor: Shiraishi, Shirou
Mitsui Toatsu Miyonodai Daini Aparto 18
2142, Tougou, Mobara-shi, Chiba-ken(JP)
Inventor: Yamada, Eichi
Miyonodai Ryo 44, 2142, Tougou
Mobara-shi, Chiba-ken(JP)
Inventor: Numata, Satoshi
954-5, 1-chome, Honda-cho
Chiba-shi, Chiba-ken(JP)**

(74) Representative: **Zumstein, Fritz, Dr. et al
Dr. F. Zumstein Dipl.-Ing. F. Klingseisen
Bräuhausstrasse 4
W-8000 München 2(DE)**

(54) **Dialkoxymethylimidazolidine derivatives, preparation thereof, insecticides containing same as an effective ingredient and intermediates therefor.**


(57) **Dialkoxymethylimidazolidine derivatives of the formula (1)**



wherein Z represents a 2-chloropyridin-5-yl group or a 2-chlorothiazol-5-yl group, R₁ and R₂, respectively, represent a lower alkyl group having from 1 to 6 carbon atoms, or a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ are joined to form a cyclic alkylene group having from 2 to 3

EP 0 455 000 A1

carbon atoms, and n is a value of 2 or 3, are described. Also, a preparation process of the derivatives and insecticidal compositions comprising the same are set forth. In addition, novel intermediate compounds of the formula (2) useful in preparing the derivatives of the formula (1) are also described along with their preparation process





15

20




35

40

45

55



wherein Z represents a 2-chloropyridin-5-yl group or a 2-chlorothiazol-5-yl group as defined before, R₁ and R₂ have, respectively, the same meanings as defined above and thus independently represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms provided that R₁ and R₂ may be joined to form a cyclic alkylene group having from 2 to 3 carbon atoms, n is a value of 2 or 3, and X represents a chlorine atom or a bromine atom.

More particularly, the compound of the formula (2) and 2-chloro-5-halomethylpyridine or 2-chloro-5-halomethylthiazole of the formula (3) are reacted in the presence of a deacidifying agent in various solvents to readily prepare the compounds of the formula (1).


Examples of the deacidifying agent include alkali metal hydroxides such as sodium hydroxide, potassium hydroxide and the like alkaline earth metal hydroxides such as magnesium hydroxide, calcium hydroxide and the like, alkali metal hydrides such as sodium hydride, potassium hydride, alkali metal alcoholates such as sodium methylate, sodium ethylate and the like, alkali metal oxides such as sodium oxide, alkali metal carbonates such as sodium carbonate, potassium carbonate and the like, alkali metal sodium hydrogencarbonates such as sodium hydrogencarbonate, potassium hydrogencarbonate and the like, hydrogensulfates such as sodium hydrogensulfate, potassium hydrogensulfate and the like, phosphates, trisodium phosphate, disodium phosphate and the like, acetates such as sodium acetate, potassium acetate and the like, organic salts such as triethylamine, DBU, DIMAP and the like, butyl lithium, sodium amide, and the like.

The solvents may include not only water, aromatic hydrocarbons such as benzene, toluene, xylene and the like, aliphatic hydrocarbons such as hexane, heptane, petroleum benzene and the like, aprotic non-polar solvents such as dimethylformamide, dimethylacetamide, dimethylsulfoxide, 1,3-dimethyl-2-imidazolidinone, 1-methyl-2-pyrrolidinone and the like, ethers such as ethyl ether, diisopropyl ether, 1,2-dimethoxyethane, tetrahydrofuran, dioxane and the like, nitriles such as acetonitrile, propionitrile and the like, and ketones such as acetone, diisopropyl ketone and the like.


When there are used phase transfer catalysts such as tetrabutylammonium bromide, triethylbenzylammonium chloride and the like, intended imidazolidine derivatives (1) can be obtained in high yield.

The reaction temperature and the reaction time can be varied over wide ranges. In general, the reaction temperature is in the range of -20 to 200°C, preferably from 0 to 100°C and the reaction time is in the range of from 0.01 to 30 hours, preferably from 0.1 to 15 hours.

The compounds (1) of the invention may include isomers of the following formula.



In the above reaction formula, the starting material of the formula (2) can be prepared according to the following reaction sequence



wherein R represents a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms, a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, R₁ and R₂ independently represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms, or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ may be joined to complete an alkylene group having from 2 to 3.

More particularly, the compounds (1) can be readily prepared in high yield by reaction between the nitroguanidine derivative of the formula (4) (literature on its preparation: J. Am. Chem. Soc., 70, 430 (1948)) and an orthoformic acid ester derivative (literature on its preparation: Synthesis, 153(1974)). The compound represented by the formula (2) is a novel compound which was synthesized by us for the first time and its synthetic reaction is a novel reaction which we first found.

The reaction is feasible in the absence of or in solvents. Examples of the solvent include aromatic hydrocarbons such as benzene, toluene, xylene and the like, halogenated hydrocarbons such as chloroform, 1,2-dichloroethane and the like, ethers such as diisopropyl ether, 1,2-dimethoxyethane, tetrahydrofuran, dioxane and the like, aprotic polar solvents such as dimethylformamide, dimethylsulfoxide, sulforan, 1,3-dimethyl-2-imidazolidinone and the like, nitriles such as acetonitrile, propionitrile and the like, and ketones such as acetone, methyl isobutyl ketone and the like.

The reaction temperature and the reaction time can be widely varied. In general, the reaction temperature is in the range of from 50 to 300°C, preferably from 70 to 200°C. The reaction is usually carried out under normal pressure conditions and may be likewise performed under pressure.


Although not necessarily required, catalysts may be used including mineral acids such as sulfuric acid, hydrochloric acid, phosphoric acid, nitric acid and the like, sulfonic acids such as p-toluenesulfonic acid, methanesulfonic acid, trifluoromethanesulfonic acid and the like, carboxylic acids such as formic acid, acetic acid, benzoic acid and the like, Lewis acids such as aluminium chloride, tin tetrachloride, zinc chloride, boron trifluoride, titanium tetrachloride and the like, pyridine hydrochloric acid salt, ammonium salts such as tetrabutylammonium chloride, and acidic or basic oxides such as zirconium oxide, silica gel, alumina and the like.

The reaction time is in the range of from 0.1 to 30 hours, preferably from 0.5 to 20 hours.

The amount of the orthoformic acid ester derivative (5) may be not less than 1.0 mole per mole of the nitroguanidine derivative (4) and is preferably in the range of from 1.0 to 10.0 moles in view of the economy.

This reaction is usually effected under normal pressure conditions and may be likewise performed under pressure.

The compound (2) of the invention may include E and Z isomers along with a tautomer as shown below



On the other hand, the chloropyridylmethyl halides of the formula (3) are known compounds and can be

prepared according to a procedure described in literature (literature on the preparation: J. Heterocyclic Chem., 16, 333 (1979) and J. Med. Chem., 14, 557 (1971)). The thiazolymethyl halides can be obtained by subjecting 2-amino-5-alkoxycarbonylthiazoles to diazotization, introducing halogen atoms, reducing the halogenated products with a lithium aluminium hydride by a usual manner, and subjecting the resulting 2-halogeno-5-hydroxymethylthiazoles to conversion with halogen atoms by a usual manner.

The derivatives of the formula (1) according to the invention have great insecticidal activity and can be used as an insecticide. The derivatives of the formula (1) of the invention show a high control effect on harmful insects without involving any phytotoxicity to cultivated plants.

Insect pests to which the derivatives of the invention can be applied, for instance, include:

10

Lepidoptera

Pieris rapae crucivora Boisduval - Common cabbageworm

Spodoptera litura Fabricius - Common cutworm

15 *Ostrinia furnacalis* Guenee - Oriental corn borer

Plutella xylostella Linne - Diamond backmoth

Chilo suppressalis Walker - Rice stem borer

Hemiptera

20

Nephotettix cincticeps Uhler - Green rice leafhopper

Nilaparvata lugens Stal - Brown rice planthopper

Laodelphax striatellus Fallen - Small brown planthopper

Unaspis yanonensis Kuwana - Arrowhead scale

25 *Myzus persicae* Sulzer - Green peach aphid

Aphis gossypii Glover - Cotton aphid

Lipaphis pseudobrassicae Davis - Turnip aphid

Nezara antennata Scott - Common green stink bug

Trialeurodes vaporariorum Westwood - Greenhouse whitefly

30

Coleoptera

Callosobruchus chinensis Linne - Azuki bean weevil

Sitophilus oryzae Linne - Rice Weevil

35 *Henosepilachna vigintioctopunctata* Fabricius - 28-spotted lady beetle

Anomala rufocuprea Motschulsky - Soy bean beetle

Leptinotarsa decemlineata Say - Colorado potato beetle

Lissorhoptrus oryzophilus Kuschel - Rice water weevil

Orthoptera

Blattella germanica Linne - German cockroach

Periplaneta americana Linne - American cockroach

Gryllotalpa africana palisot de Beauvois - African mole cricket

45 *Locusta migratoria danica* Linne - Asiatic locust

Reticulitermes speratus Kolbe

Coptotermes formosanus Shiraki - Formosan subterranean termite

Diptera

50

Musca domestica vicina Macquart - House fly

Aedes aegypti Linne - Yellow fever mosquito

Culex pipiens pallens Coquillett

Culex tritaeniorhynchus Giles

55 Where the compounds of the formula (1) of the invention is actually applied, it may be used singly without addition of any other ingredient. However, it is usual to formulate carriers in order to make easy application as a control chemical.

For preparation of the compounds of the invention, any specific requirement is not necessary and

optional preparations, such as emulsions, dusts, granules, fine powders, oils, aerosols, poisonous feeds and the like, according to the procedures of preparing general agricultural chemicals well known in the art.

The term "carrier" used herein is intended to mean synthetic or natural, organic or inorganic materials which assist the effective ingredient to arrive at sites or portions to be treated and which are formulated in order to make easy storage, transport and handling of the effective compound. Appropriate solid carriers include, for example, clays such as montmorillonite, kaolin and the like, inorganic substances such as diatomaceous earth, white clay, talc, vermiculite, gypsum, calcium carbonate, silica gel, ammonium sulfate and the like, plant organic substances such as soybean flour, saw dust, wheat flour and the like, and urea.

Suitable liquid carriers include, for example, aromatic hydrocarbons such as toluene, xylene, cumene and the like, paraffin hydrocarbons such as kerosene, mineral oils and the like, halogenated hydrocarbons such as carbon tetrachloride, chloroform, dichloroethane and the like, ketones such as acetone, methyl ethyl ketone and the like, ethers such as dioxane, tetrahydrofuran and the like, alcohols such as methanol, ethanol, propanol, ethylene glycol and the like, dimethylformamide, dimethyl sulfoxide, water and the like.

In order to reinforce the efficacy of the compound of the formula (1) of the invention, the following adjuvants may be used singly or in combination, depending on the type of preparation, the manner of application and the purpose.

For the purposes of emulsification, dispersion, spreading, wetting, bonding and stabilization, there are used water-soluble salts such as ligninsulfonates, nonionic surface active agents such as alkylbenzene sulfonates, alkylsulfates and the like, lubricants such as calcium stearate, waxes and the like, stabilizers such as isopropoxyhydrogenphosphates, and methyl cellulose, carboxymethyl cellulose, casein, gum arabi and the like. It should be noted that the adjuvants are not limited to those mentioned above and other adjuvants ordinarily used for this purpose may also be used.

The compounds of the formula (1) of the invention may develop better insecticidal activity when used in combination of two or more. If other physiologically active substances or chemicals are used in combination, multi-purpose compositions with good efficacy can be prepared with the possibility of developing a synergistic effect. Examples of such physiologically active substances include: synthetic pyrethroids, and isomers thereof or pyrethrum extracts, such as allethrin, N-(chrysanthemoylmethyl)-3,4,5,6-tetrahydrophthalimide, 5-benzyl-3-furylmethyl chrysanthemate, 3-phenoxybenzyl chrysanthemate, 5-propargylfurfuryl chrysanthemate and other known cyclopropanecarboxylic acid esters, 3-phenoxybenzyl 2,2-dimethyl-3-(2,2-dichlorovinyl)-cyclopropane-1-carboxylate, 2,2-dimethyl-3-(2,2-dibromovinyl)-cyclopropane-1-carboxylate, 3-phenoxy- α -cyanobenzyl α -isopropyl-4-chlorophenylacetate and the like; organophosphate insecticides such as O,O-diethyl-O-(3-oxo-2-phenyl-2H-pyridazin-6-yl)phosphorothioate (available from Mitsui-Toatsu Chem. Ind. Co., Ltd. under the trade name of Ofunack), O,O-dimethyl-O-(2,2-dichlorovinyl)phosphate (DDVP), O,O-dimethyl-O-(3-methyl-4-nitrophenyl)phosphorothioate, diazinon, O,O-dimethyl-O-4-cyanophenylphosphorothioate, O,O-dimethyl-S-[α -(ethoxycarbonyl)benzyl]phosphorodithioate, 2-methoxy-4H-1,3,2-benzodioxaphosphorin-2-sulfide, O-ethyl-O-4-cyanophenylphosphonothioate and the like; carbamate insecticides such as 1-naphthyl N-methylcarbamate (NAC), m-tolyl N-methylcarbamate (MTMC), 2-dimethylamino-5,6-dimethylpyrimidin-4-yl dimethylcarbamate (Pyrimet), 3,4-dimethylphenyl N-methylcarbamate, 2-isopropoxyphenyl N-methylcarbamate and the like; aryl propyl ether insecticides such as 3-phenoxybenzyl 2-(4-chlorophenyl)-2-methyl propyl ether, 3-phenoxy-4-fluorobenzyl 2-(4-chlorophenyl)-2-methylpropyl ether, 3-phenoxybenzyl 2-(4-ethoxyphenyl)-2-methylpropyl ether, 3-phenoxy-4-(fluorobenzyl 2-(4-ethoxyphenyl)-2-methylpropyl ether and the like; aromatic alkane insecticides such as 1-(3-phenoxyphenyl)-4-(4-chlorophenyl)-4-methylpentane, 1-(3-phenoxy-4-fluorophenyl)-4-(4-chlorophenyl)-4-methylpentane, 1-(3-phenoxyphenyl)-4-(4-ethoxyphenyl)-4-methylpentane, 1-(3-phenoxy-4-fluorophenyl)-4-(4-ethoxyphenyl)-4-methylpentane and the like; and other insecticides, acaricides, fungicides, nematocides, herbicides, plant growth regulators, fertilizers, BT agents, insect hormone compounds, and other agricultural chemicals.

Although the compounds of the formula (1) of the invention are stable against light, heat and oxidation, antioxidants or UV absorbers may be added in appropriate amounts, if necessary, including, for example, phenol derivatives or bisphenol derivatives such as BHT (2,6-di-t-butyl-4-methylphenol), BHA (butylhydroxyanisole) and the like, arylamines or benzophenone compounds such as phenyl- α -naphthylamine, phenyl- β -naphthylamine, a condensate of phenetidine and acetone, thereby obtaining more stable compositions.

When the compounds of the formula (1) of the invention are used as an insecticide, they are used in an amount of from 0.0001 to 95 wt%, preferably from 0.01 to 50 wt% of the insecticide.

When the insecticide of the invention is applied, the effective ingredient is used at a concentration of 0.01 to 5000 ppm, preferably from 0.1 to 1000 ppm.

The application amount per 10 ares is generally in the range of from 1 to 300 g of the effective

ingredient.

The present invention is more particularly described by way of examples, which should not be construed as limiting the invention.

5 Synthesis Example 1 (Compound No. 1)

17.6 g of 1-diethoxymethyl-2-nitroiminoimidazolidine and 21.0 g of anhydrous potassium carbonate were added to 200 ml of dimethylformamide. While agitating at 70 °C, a solution of 12.3 g of 2-chloro-5-chloromethylpyridine in 30 ml of dimethylformamide was dropped in the mixture.

10 After completion of the dropping, the reaction mixture was poured into water, followed by extraction with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure to give 25.0 g of an oily residue. This was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to obtain 11.3 g of 1-diethoxymethyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

15 Synthesis Example 2 (Compound No. 3)

15.0 g of 1-(1-ethoxy-1-propoxy)methyl-2-nitroiminoimidazolidine, 11.6 g of anhydrous potassium carbonate, 11.3 g of 2-chloro-5-chloromethylpyridine and 60 ml of dimethyl sulfoxide were agitated at 75 °C for 1 hour. The reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous sodium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oily residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 4.7 g of 1-(1-ethoxy-1-propoxy)methyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

25 Synthesis Example 3 (Compound No. 5)

6.8 g of 1-[1-ethoxy-1-(2-methoxyethoxy)]methyl-2-nitroiminoimidazolidine, 4.8 g of anhydrous potassium carbonate, 6.0 g of 2-chloro-5-chloromethylpyridine and 20 ml of dimethyl sulfoxide were agitated at 70 °C for 1 hour.

30 The reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous sodium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oily residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 2.8 g of 1-[1-ethoxy-1-(2-methoxyethoxy)]methyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

Synthesis Example 4 (Compound No. 6)

16.0 g of 1-[bis-(2-chloroethoxy)methyl]-2-nitroiminoimidazolidine, 13.0 g of anhydrous potassium carbonate, 13.0 g of 2-chloro-5-chloromethylpyridine and 70 ml of dimethyl sulfoxide were agitated at 70 °C for 1 hour. The reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous sodium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oily residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 14.0 g of 1-[bis-(2-chloroethoxy)methyl]-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

Synthesis Example 5 (Compound No. 7)

7.0 g of 1-[1-ethoxy-1-(2,2,2-trifluoroethoxy)]methyl-2-nitroiminoimidazolidine, 7.0 g of anhydrous potassium carbonate, 7.0 g of 2-chloro-5-chloromethylpyridine and 45 ml of dimethyl sulfoxide were agitated at 70 °C for 1 hour.

50 The reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous sodium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oily residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 2.3 g of 1-[1-ethoxy-1-(2,2,2-trifluoroethoxy)]methyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

Synthesis Example 6 (Compound No. 9)

6.0 g of 1-dimethoxymethyl-2-nitroiminoimidazolidine and 8.6 g of anhydrous potassium carbonate were added to 50 ml of dimethyl sulfoxide. While agitating at 70 °C, a solution of 5.5 g of 2-chloro-5-chloromethylpyridine in 10 ml of dimethyl sulfoxide was dropped in the mixture in 20 minutes, followed by further agitation at the same temperature for 30 minutes.

- 5 After completion of the dropping, the reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure to give 13.7 g of an oily residue. The residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 4.8 g of 1-dimethoxymethyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)imidazolidine.

10

Synthesis Example 7 (Compound No. 10)

- 5.5 g of 1-diethoxymethyl-2-nitroiminohexahydropyrimidine and 6.5 g of anhydrous potassium carbonate were added to 50 ml of dimethylformamide. While agitating at 70 °C, a solution of 3.8 g of 2-chloro-5-chloromethylpyridine in 10 ml of dimethylformamide was dropped in the mixture, followed by further agitation at the same temperature for 40 minutes.

- After completion of the dropping, the reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure to give 11.0 g of an oily residue. The residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 2.3 g of 1-diethoxymethyl-2-nitroimino-3-(2-chloropyridin-5-ylmethyl)hexahydropyrimidine.

20

Synthesis Example 8 (Compound No. 11)

- 25 5.0 g of 1-diethoxymethyl-2-nitroiminoimidazolidine and 6.0 g of anhydrous potassium carbonate were added to 50 ml of dimethyl sulfoxide. While agitating at 70 °C, a solution of 3.9 g of 2-chloro-5-chloromethylthiazole in 5 ml of dimethyl sulfoxide was dropped in the mixture, followed by further agitation at the same temperature for 1.5 hours.

- After completion of the dropping, the reaction mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate and the solvent was distilled off under reduced pressure to give 9.8 g of an oily residue. The residue was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 5.4 g of 1-diethoxymethyl-2-nitroimino-3-(2-chlorothiazol-5-ylmethyl)imidazolidine.

- Typical compounds of the formula (1) which were prepared according to procedures similar to those described in Examples 1 to 8 are shown in Tables 1 and 2.

35


40

45

50

55

Table 1



(Z = 2-chloropyridin-5-yl group)

Compound No.	Substituents			Values of Physical Properties															
	R ₁	R ₂	n																
1	Et	Et	2	<p>δ_{TMS} (CDCl₃) (ppm) : 1.25(6H,t, J=6.9Hz), 3.52~3.82(8H,m), 4.48(2H,s), 5.76(1H,s), 7.37(1H,d, J_{AB}=8.4Hz), 7.72(1H,dd, J_{AB}=8.4Hz, J=2.5Hz), 8.32(1H,d, J=2.5Hz)</p> <p>ν_{MAX} (KBr) (cm⁻¹) : 1560, 1520, 1460, 1425, 1250, 1100, 1050</p> <p>Elementary Analysis (C₁₄H₂₀ClN₅O₄) :</p> <table> <tr> <td></td><td>C</td><td>H</td><td>Cl</td><td>N</td></tr> <tr> <td>Calculated (%)</td><td>47.00</td><td>5.63</td><td>9.91</td><td>19.57</td></tr> <tr> <td>Found (%)</td><td>47.52</td><td>5.61</td><td>9.72</td><td>20.22</td></tr> </table> <p>m.p. : 87.0~87.5°C</p>		C	H	Cl	N	Calculated (%)	47.00	5.63	9.91	19.57	Found (%)	47.52	5.61	9.72	20.22
	C	H	Cl	N															
Calculated (%)	47.00	5.63	9.91	19.57															
Found (%)	47.52	5.61	9.72	20.22															
2	n-Pr	n-Pr	2	<p>δ_{TMS} (acetone-d₆) (ppm) : 0.92~0.99(6H,m), 1.57~1.67(4H,m), 3.44~3.51(2H,m), 3.53~3.61(4H,m), 3.76~3.84(2H,m), 4.48(2H,s), 5.75(1H,s), 7.36(1H,d, J=8.1Hz), 7.72(1H,dd, J=8.1Hz, J=2.2Hz), 8.33(1H,d, J=2.2Hz)</p> <p>ν_{MAX} (neat) (cm⁻¹) : 1552, 1462, 1255, 1101</p>															

Table 1 (continued)

Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
3	nPr	Et	2	δ_{TMS} (acetone-d ₆) (ppm): 0.92 ~ 0.99 (3H,m), 1.21 ~ 1.28 (3H,m), 1.57 ~ 1.69 (2H,m), 3.44 ~ 3.61 (6H,m), 3.74 ~ 3.84 (2H,m), 4.54 (2H,s), 5.75 (1H,s), 7.34 ~ 7.38 (1H,m), 7.69 ~ 7.73 (1H,m), 8.32 (1H,s) ν_{MAX} (neat) (cm ⁻¹): 1557, 1455, 1100
4	iso-Pr	iso-Pr	2	δ_{TMS} (acetone-d ₆) (ppm): 1.15 ~ 1.20 (12H,m), 3.78 ~ 3.93 (6H,m), 4.55 (2H,s), 5.86 (1H,s), 7.47 (1H,d, J=8.8Hz), 7.85 (1H,dd, J=8.8Hz, J=2.2Hz), 8.40 (1H,d, J=2.2Hz) ν_{MAX} (KBr) (cm ⁻¹): 1564, 1255, 1100 m.p.: 78 ~ 79°C
5	CH ₃ OC ₂ H ₄	Et	2	δ_{TMS} (acetone-d ₆) (ppm): 1.19 (3H,t, J=7.2Hz), 3.30 (3H,s), 3.51 ~ 3.82 (10H,m), 4.55 (2H,s), 5.79 (1H,s), 7.47 (1H,d, J=8.1Hz), 7.86 (1H,dd, J=8.1Hz, J=2.2Hz), 8.42 (1H,d, J=2.2Hz) ν_{MAX} (neat) (cm ⁻¹): 1553, 1461, 1257, 1106
6	ClC ₂ H ₄	ClC ₂ H ₄	2	δ_{TMS} (acetone-d ₆) (ppm): 3.75 ~ 4.00 (12H,m), 4.57 (2H,s), 5.92 (1H,s), 7.46 (1H,d, J=7.1Hz), 7.85 (1H,dd, J=7.1Hz, J=2.2Hz), 8.41 (1H,d, J=2.2Hz) ν_{MAX} (neat) (cm ⁻¹): 1558, 1463, 1255, 1106


Table 1 (continued)

Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
7	CF ₃ CH ₂	Et	2	δ_{TMS} (acetone-d ₆) (ppm): 1.23(3H,m), 3.6~3.77(2H,m), 3.83~3.87(4H,m), 4.21(2H,q, J=8.8Hz), 4.57(2H,s), 5.94(1H,s), 7.47(1H,d, J=8.1Hz), 7.86(1H,dd, J=8.1Hz, J=2.2Hz), 8.42(1H,d, J=2.2Hz) ν_{MAX} (neat) (cm ⁻¹): 1558, 1455, 1277, 1104
8	CF ₃ CH ₂	Me	2	δ_{TMS} (acetone-d ₆) (ppm): 3.42(3H,s), 3.6~4.00(4H,m), 4.16(2H,q, J=13.5Hz), 4.53(2H,s), 5.81(1H,s), 7.37(1H,d, J=8.1Hz), 7.67~7.90(1H,m), 8.30~8.43(1H,m) ν_{MAX} (neat) (cm ⁻¹): 1558, 1455, 1278, 1109
9	Me	Me	2	δ_{TMS} (acetone-d ₆) (ppm): 3.37(6H,s), 3.7~3.85(4H,m), 4.56(2H,s), 5.56(1H,s), 7.48(1H,d, J=8.1Hz), 7.87(1H,dd, J=8.1Hz, J=2.2Hz), 8.43(1H,d, J=2.2Hz) ν_{MAX} (KBr) (cm ⁻¹): 1561, 1535, 1447, 1282, 1099 m.p.: 62.0~64.0°C

Table 1 (continued)

Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
10	Et	Et	3	δ_{TMS} (CDCl ₃) (ppm): 1.24(6H, t, J=7.3Hz), 2.01~2.06(2H, m), 3.33(2H, t, J=5.9Hz), 3.52(2H, t, J=5.9Hz), 3.55~3.62(2H, m), 3.66~3.73(2H, m), 4.61(2H, s), 6.09(1H, s), 7.37(1H, d, J=8.3Hz), 7.76(1H, dd, J=8.3Hz, J=2.4Hz), 8.33(1H, d, J=2.2Hz) ν_{MAX} (neat) (cm ⁻¹): 1589, 1502, 1408, 1290, 1101 n_D : 1.5486(20°C)

Table 2



(Z = 2-chlorothiazol-5-yl group)

Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
11	Et	Et	2	δ_{TMS} (DMSO-d ₆) (ppm): 1.14(6H, t, J=7.3Hz), 3.47~3.61(4H, m), 3.63~3.71(4H, m), 4.52(2H, s), 5.65(1H, s), 7.68(1H, s) ν_{MAX} (neat) (cm ⁻¹): 1528, 1419, 1255, 1100

Then, preparation of intermediate compounds of the general formula (2) is described.

55 Synthesis Example 9 (Intermediate No. 1)

A mixture of 25 g of 2-nitroimidazoimidazolidin, 100 g of ethyl orthoformate and 25 ml of 1,3-dimethyl-2-imidazolidinon was heated under reflux for 3 hours. After cooling to room temperature, the mixture was

poured into water, followed by extraction with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate, after which the solvent was distilled off under reduced pressure. The resultant crystals were sludged with ether to give 32 g of 1-diethoxymethyl-2-nitroiminoimidazolidine.

5 Synthesis Example 10 (Intermediate No. 5)

16.5 g of 2-nitroiminoimidazolidine and 18.0 g of ethylenedioxyethyl ether were heated under reflux for about 4 hours while removing the resultant distillate by means of the Dean-Stark trap. After cooling to room temperature, the reaction mixture was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 2.8 g of 1-di-n-propoxymethyl-2-nitroiminoimidazolidine.

Synthesis Example 11 (Intermediate No. 6)

9.9 g of 2-nitroiminoimidazolidine and 18.0 g of n-propyl orthoformate were heated under reflux for about 4 hours while removing a distillate by means of Dean-Stark trap. After cooling to room temperature, the reaction mixture was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 1.9 g of 1-di-n-propoxymethyl-2-nitroiminoimidazolidine.

Synthesis Example 12 (Intermediate No. 9)

22.0 g of methyl orthoformate was dropped in about 2 hours in a mixture of 10 g of 2-nitroiminoimidazolidine, 20 ml of 1,3-dimethyl-2-imidazolidinone and 0.05 g of sulfuric acid at 150 °C, followed by heating under reflux for further 1 hour while removing the resultant distillate by means of the Dean-Stark trap. After cooling to room temperature, the mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate, after which the solvent was distilled off under reduced pressure. The resultant crude crystals were sludged with ether to give 1.9 g of 1-dimethoxymethyl-2-nitroiminoimidazolidine.

Synthesis Example 13 (Intermediate No. 10)

20.0 g of methyl orthoformate was dropped in about 1.5 hours in a mixture of 10 g of 2-nitroiminohexahydropyrimidine, 20 ml of 1,3-dimethyl-2-imidazolidinone and 0.05 g of sulfuric acid at 150 °C, followed by heating under reflux for further 1 hour while removing the resultant distillate by means of the Dean-Stark trap. After cooling to room temperature, the mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oily substance was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to give 1.5 g of 1-dimethoxymethyl-2-nitroiminohexahydropyrimidine.

40 Synthesis Example 14 (Intermediate No. 11)

A mixture of 10 g of 2-nitroiminohexahydropyrimidine, 12.4 g of ethyl orthoformate, 10 ml of 1,3-dimethyl-2-imidazolidinone was heated under reflux for 2.0 hours while removing the resultant distillate by means of the Dean-Stark trap. After cooling to room temperature, the mixture was poured into water, and extracted with ethyl acetate. After washing with water, the extract was dried with anhydrous magnesium sulfate, after which the solvent was distilled off under reduced pressure. The resultant oil was purified by column chromatography [silica gel, eluent: hexane/ethyl acetate (1:2)] to obtain 2.3 g of 1-diethoxymethyl-2-nitroiminohexahydropyrimidine. Typical examples of the compounds of the formula (2) which could be prepared according to the procedures similar to those of Examples 9 to 14 are shown in Table 3.




Table 3

Intermediate Compound No.	Substituents			Values of Physical Properties												
	R ₁	R ₂	n													
1	Et	Et	2	δ_{TMS} (CDCl ₃) (ppm) : 1.24 (6H, t, J=6.9Hz), 3.44 ~ 3.87 (8H, m), 5.95 (1H, s), 8.36 (1H, s) ν_{MAX} (KBr) (cm ⁻¹) : 3340, 1570, 1530, 1470, 1440, 1280, 1220, 1170, 1090, 1040 Elementary Analysis (C ₈ H ₁₆ N ₄ O ₄) : <table><tr><td></td><td>C</td><td>H</td><td>N</td></tr><tr><td>Calculated (%)</td><td>41.37</td><td>6.94</td><td>24.13</td></tr><tr><td>Found (%)</td><td>40.94</td><td>6.90</td><td>24.58</td></tr></table> m.p. : 100.2 ~ 101.8 °C		C	H	N	Calculated (%)	41.37	6.94	24.13	Found (%)	40.94	6.90	24.58
	C	H	N													
Calculated (%)	41.37	6.94	24.13													
Found (%)	40.94	6.90	24.58													
2	CH ₃ OC ₂ H ₄	CH ₃ OC ₂ H ₄	2	δ_{TMS} (acetone-d ₆) (ppm) : 3.33 (6H, s), 3.45 ~ 3.87 (12H, m), 5.98 (1H, s), 8.30 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹) : 3375, 1575, 1447, 1108 m.p. : 68.0 ~ 70.5 °C												
3	ClC ₂ H ₄	ClC ₂ H ₄	2	δ_{TMS} (acetone-d ₆) (ppm) : 3.71 ~ 4.01 (12H, m), 6.04 (1H, s), 8.50 ~ 8.90 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹) : 3359, 1574, 1443, 1290, 1098 m.p. : 85.0 ~ 89.5 °C												

Table 3 (continued)

Intermediate Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
4	iso-Pr	iso-Pr	2	δ_{TMS} (acetone-d ₆) (ppm): 1.16 ~ 1.21 (12H, m), 3.66 ~ 3.92 (6H, m), 6.01 (1H, s), 8.71 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹): 3423, 1565, 1433, 1283, 1083 m.p.: 137 ~ 146°C
5	CH ₂ CH ₂		2	δ_{TMS} (acetone-d ₆) (ppm): 3.55 ~ 4.10 (8H, m), 6.49 (1H, s), 8.08 ~ 8.45 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹): 3359, 1587, 1444, 1302, 1103 m.p.: 123 ~ 126.5°C
6	nPr	nPr	2	δ_{TMS} (CDCl ₃) (ppm): 0.92 ~ 0.99 (6H, m), 1.56 ~ 1.71 (4H, m), 3.42 ~ 3.62 (4H, m), 3.69 ~ 3.74 (2H, m), 3.81 ~ 3.85 (2H, m), 5.94 (1H, s), 8.38 (1H, broad-s) ν_{MAX} (neat) (cm ⁻¹): 3416, 1574, 1447, 1288, 1100
7	CF ₃ CH ₂	Et	2	δ_{TMS} (acetone-d ₆) (ppm): 1.23 (3H, t, J=6.6Hz), 3.64 ~ 3.90 (6H, m), 4.20 (2H, q, J=8.8Hz), 6.06 (1H, s), 8.84 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹): 3384, 1574, 1447, 1297, 1090 m.p.: 80 ~ 85°C
8	CF ₃ CH ₂	Me	2	δ_{TMS} (acetone-d ₆) (ppm): 3.42 (3H, s), 3.59 ~ 4.22 (6H, m), 5.92 (1H, s), 8.10 ~ 8.52 (1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹): 3397, 1582, 1444, 1286, 1110 m.p.: 144°C (dec.)

Table 3 (continued)

Intermediate Compound No.	Substituents			Values of Physical Properties
	R ₁	R ₂	n	
9	Me	Me	2	δ_{TMS} (CDCl ₃) (ppm) : 3.41(6H,s), 3.66~3.71(2H,m), 3.81~3.86(2H,m), 5.78(1H,s), 8.40(1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹) : 3367, 1577, 1533, 1289, 1104 m.p. : 118~125°C
10	Me	Me	3	δ_{TMS} (CDCl ₃) (ppm) : 2.04~2.09(2H,m), 3.40(6H,s), 3.45~3.70(4H,m), 6.28(1H,s), 9.80(1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹) : 3284, 1548, 1423, 1236, 1099
11	Et	Et	3	δ_{TMS} (CDCl ₃) (ppm) : 1.23(6H,t, J=7.8Hz), 1.98~2.03(2H,m), 3.48(4H,q, J=7.8Hz), 3.51~3.57(2H,m), 3.64~3.70(2H,m), 6.50(1H,s), 9.98(1H, broad-s) ν_{MAX} (KBr) (cm ⁻¹) : 3284, 1544, 1427, 1234, 1097 m.p. : 85.5~86.5°C

The compositions of the invention are more particularly described by way of Preparation Examples

Formulation Example 1

20 parts by weight of the compound of the invention prepared in Synthesis Example 1, 10 parts by

weight of Sorpol 355S (surfactant available from Toho Chem. Co., Ltd.) and 70 parts by weight of xylene were uniformly agitated and mixed to give an emulsifiable concentrate.

Formulation Example 2

20 parts by weight of the compound of the invention prepared in Synthesis Example 1, 2 parts by weight of sodium alkyl naphthalenesulfonate, 5 parts by weight of sodium ligninsulfonate, 5 parts by weight of white carbon and 68 parts by weight of diatomaceous earth were uniformly agitated and mixed to give a wettable powder.

Formulation Example 3

0.3 parts by weight of the compound of the invention prepared in Synthesis Example 1 was dissolved in acetone. While mixing with 99.7 parts by weight of clay, the acetone was evaporated to give a powder.

Formulation Example 4

2 parts by weight of the compound of the invention prepared in Synthesis Example 1, 2 parts by weight of sodium ligninsulfonate, and 96 parts by weight of bentonite were uniformly divided into pieces and mixed, to which water was added for kneading, followed by granulation and drying to give a granular.

The insecticidal activity of the compounds of the formula (1) is particularly by way of test examples.

Test Example 1 Effect on *Laodelphax striatellus* Fallén - smaller brown planthopper

The emulsion prepared in Formulation Example 1 was diluted to predetermined concentrations and 2 ml of each diluted emulsion was applied over a bundle of several rice seedlings (about third leaf stage). After drying in air, the treated seedlings were covered with a metal gauze cylinder, in which ten female adults of the smaller brown planthopper were released, followed by placing in a temperature controlled room at 25°C. After 48 hours, the mortality was checked. The results are shown in Table 4. For a control chemical, there was used the compound 1-methoxymethyl-2-nitroimino 3-(2-chloropyridin-5-ylmethyl)imidazolidine as shown below.




Table 4

Test Compound No.	Mortality (%)	
	100 ppm	10 ppm
No. 1	100	100
No. 2	100	100
No. 3	100	100
No. 4	100	100
No. 5	100	95
No. 6	100	100
No. 7	100	100
No. 8	100	90
No. 9	100	100
No. 10	100	100
No. 11	100	100
Reference Compound	30	0
Non treated	0	0

Test Example 2 Effect on resistant strain of *Nephotettix cincticeps* Uhler - resistant green rice leafhopper

The emulsion prepared in Formulation Example 1 was diluted to predetermined concentrations and each solution was applied in an amount of 3 ml over a bundle of several rice seedlings (about third leaf stage). After drying in air, the treated seedlings were covered with a metal gauze cylinder, in which ten female adults of leafhopper that is resistant to organophosphate and carbamate agents were released, followed by placing in a temperature controlled room. After 48 hours, the mortality was checked. The results are shown in Table 5. For a control chemical, there was used the same compound in Test Example 1.

Table 5

Test Compound No.	Mortality (%)	
	10 ppm	1 ppm
No. 1	100	100
No. 2	100	100
No. 3	100	100
No. 4	100	100
No. 5	100	100
No. 6	100	90
No. 7	100	100
No. 8	100	100
No. 9	100	100
No. 10	100	100
No. 11	100	100
Reference Compound	100	45
Non treated	0	0

Test Example 3 Effect on *Callosobruchus chinensis* Linne - Azuki bean weevil

An acetone solution of the compound of the invention prepared in Synthesis Example 1 was added to Petri dish with a diameter of 9 cm, followed by removal of the acetone by evaporation. Twenty female adults of the Azuki bean weevil, which were 2 to 3 days after emergence were placed in the dish at 25°C. After 48 hours, the mortality was checked. The results are shown in Table 6. For a control chemical, there was used diazinon of the formula (7) [O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].

Table 6

Test Compound No.	Mortality (%)	
	0.1 mg/dish	0.001 mg/dish
No. 1	100	100
Diazinone	95	30
Non treated	2.5	

Test Example 4 Effect on *Myzus persicae* Sulzer - green peach aphid

Dilutions of the emulsifiable concentrate prepared in Formulation Example 1 were sprayed in an amount of 20 ml over potted eggplant seedlings (fourth or fifth leaf stage) on which the green peach aphid had been parasitic. After the application, the seedlings were placed in a greenhouse and, after three days, the number of the aphids were checked. The results are shown in Table 7. For a control chemical, there was used the same compound in Test Example 1.


Table 7

Test Compound No.	Number of Parasitic aphids	
	Concentration (1 ppm)	
	Prior to Treatment	After Three Days
No. 1	125	0
No. 2	98	0
No. 3	111	0
No. 4	106	0
No. 5	132	0
No. 6	89	0
No. 7	94	0
No. 8	131	0
No. 9	105	0
Reference Compound	120	92
Non treated	103	146

As will be apparent from the foregoing description, the imidazolidine derivatives of the formula (1) according to the invention have high insecticidal efficacy and a wide insecticidal spectrum. The imidazolidine derivatives of the formula (1) can be readily prepared according to a process of the invention using novel intermediates of the formula (2). The agricultural chemicals comprising the imidazolidine derivatives of the formula (1) have good characteristics as an insecticide.

Claims


1. An imidazolidine derivative of the formula (1)




wherein Z represents a 2-chloropyridin-5-yl group or a 2-chlorothiazol-5-yl group, R₁ and R₂, respec-

tively, represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ are joined to form a cyclic alkylene group having from 2 to 3 carbon atoms, and n is a value of 2 or 3.


- 5
2. An imidazolidine derivative according to Claim 1, wherein, in the formula (1), R₁ and R₂ are independently a lower alkyl group having from 1 to 4 carbon atoms and Z is a 2-chloropyridin-5-yl group.
- 10 3. A process for preparing an imidazolidine derivative of the formula (1)



25 Z represents a 2-chloropyridin-5-yl group or a 2-chlorothiazol-5-yl group, R₁ and R₂, respectively, represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ are joined to form a cyclic alkylene group having from 2 to 3 carbon atoms, and n is a value of 2 or 3, the process comprising reacting a compound of the formula (2)




wherein R₁ and R₂ independently represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ are joined to form a cyclic alkylene group having from 2 to 3 carbon atoms, and n is a value of 2 or 3, and a compound of the formula (3)




wherein Z represents a 2-chloropyridin-5-yl group or a 2-chlorothiazol-5-yl group, and X represents a chlorine atom or a bromine atom.

4. A process according to Claim 3, wherein R₁ and R₂ are independently a lower alkyl group having from 1 to 4 carbon atoms and Z is a 2-chloropyridin-5-yl group.
- 55 5. An insecticidal composition comprising an effective amount of a compound of the formula (1)

7. An imidazolidine derivative of the formula (2)




9. A process for preparing an imidazolidine derivative of the formula (2)




25

joined to form a cyclic alkylene group having from 2 to 3 carbon atoms, and n is a value of 2 or 3, the process comprising reacting a compound of the formula (4)



15 wherein n is a value of 2 or 3, and a compound of the formula (5)



20 wherein R represents a lower alkyl group having from 1 to 6 carbon atoms or a lower haloalkyl group having from 1 to 3 carbon atoms, and R₁ and R₂, respectively, represent a lower alkyl group having from 1 to 6 carbon atoms, a lower haloalkyl group having from 1 to 3 carbon atoms or a lower alkyl group having from 1 to 3 carbon atoms and substituted with a lower alkoxy group having from 1 to 3 carbon atoms, or R₁ and R₂ are joined to form a cyclic alkylene group having from 2 to 3 carbon atoms.

- 25
10. A process according to Claim 9, wherein R, R₁ and R₂ independently represent a lower alkyl group having from 1 to 4 carbon atoms.



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 91 10 5171

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 315 826 (NIHON TOKUSHU NOYAKU SEIZO K.K.) - - -		C 07 D 401/06 A 01 N 43/50
A	EP-A-0 285 985 (BAYER AG) - - -		A 01 N 43/54 A 01 N 43/78
A	PATENT ABSTRACTS OF JAPAN, vol. 12, no. 426 (C-542)[3273], 10th November 1988; & JP-A-63 156 786 (NIPPON TOKUSHU NOYAKU SEIZO) 29-06-1988 * The whole document *		A 01 N 43/40 C 07 D 417/06 C 07 D 405/14 C 07 D 417/14
A	PATENT ABSTRACTS OF JAPAN, vol. 13, no. 362 (C-625)[3710], 14th August 1989; & JP-A-01 121 287 (NIPPON TOKUSHU NOYAKU SEIZO K.K.) 12-05-1989 * The whole document *		
D,A	PATENT ABSTRACTS OF JAPAN, vol. 11, no. 282 (C-446)[2729], 11th September 1987; & JP-A-62 81 382 (NIPPON TOKUSHU NOYAKU SEIZO K.K.) 14-04-1987 * The whole document *		
D,A	EP-A-0 277 317 (NIHON TOKUSHU NOYAKU SEIZO K.K.) - - - - -		
The present search report has been drawn up for all claims			
Place of search The Hague		Date of completion of search 11 July 91	Examiner DE BUYSER I.A.F.
<div>CATEGORY OF CITED DOCUMENTS</div> <div>X: particularly relevant if taken alone Y: particularly relevant if combined with another document of the same category A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention</div> <div>E: earlier patent document, but published on, or after the filing date D: document cited in the application L: document cited for other reasons &: member of the same patent family, corresponding document</div>			